

# Betacol®

1375

Please Copy for Your Patients

## Betacol Contains Betaine, Vitamin B<sub>6</sub>, Niacin, and Phytochemicals to Promote Healthy Liver, Cardiovascular, and Cellular Function

Betacol contains *Tillandsia usneoides*, a source of important phytochemicals that contribute to cardiovascular and cellular health. Several bioactive compounds, including vitamin C, carotene, and HMG (3-hydroxy-3-methylglutaric acid), are found in *Tillandsia usneoides*. Vitamins, minerals, and phytochemicals in Betacol work at the cellular level to maintain healthy blood glucose and cholesterol levels in individuals with normal levels. Betaine in Betacol is important for healthy liver function, especially the metabolism of fat.†

### How Betacol Keeps You Healthy

#### Promotes healthy liver function

Betaine is a lipotropic agent that promotes transportation and metabolism of fats, helping to prevent the accumulation of fat in the liver. Several studies demonstrate betaine's unique ability to support a healthy liver.†

#### Supports cardiovascular health

*Tillandsia usneoides* supports arterial health by helping to maintain normal serum cholesterol levels in individuals with normal levels. HMG (3-hydroxy-3-methylglutaric acid), the bioactive compound in *Tillandsia usneoides*, has been shown in several studies to help support cardiovascular health factors. Several studies have confirmed that HMG operates at the enzymatic level in the kidney and intestine. Studies also indicate that HMG may help maintain healthy cardiovascular function. Betacol contains vitamin B<sub>6</sub>, which is important for red blood cell formation and helps maintain optimum homocysteine levels in the blood. Vitamin B<sub>6</sub> is well documented for its ability to support the heart muscle and associated blood vessels.†

#### Promotes healthy cellular function

Vitamin B<sub>6</sub> is required to synthesize the nucleic acids RNA and DNA—the molecules that carry the genetic instructions for normal cellular growth and reproduction. Betacol also contains niacin, which functions as a coenzyme that is key to cellular respiration, carbohydrate and protein metabolism, and lipid synthesis. Niacin is a hydrogen acceptor, combining with hydrogen atoms as they are removed from the food we eat. Once the hydrogen atoms are removed from the food, the coenzyme continues transferring them to other compounds in a series of complex oxidation reactions. Several studies indicate that extracts from *Tillandsia usneoides* contain compounds capable of reducing serum glucose. HMG may also help support liver function.†



**Introduced in:**

1946

**Content:**

40 Capsules

#### Supplement Facts:

Serving Size: 1 capsule  
Servings per Container: 40

		%DV
Calories	2	
Niacin	10 mg	50%
Vitamin B <sub>6</sub>	2 mg	100%

Betacol® 1375



800-558-8740 • [www.standardprocess.com](http://www.standardprocess.com)

† These statements have not been evaluated by the Food & Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

## What Makes Betacol Unique

### Unique Product Attributes

#### A distinctive product that contains choline

- Choline is essential to metabolize fat cholesterol, proteins, and carbohydrates effectively†

#### Multiple nutrients from a variety of plant and animal sources

- Extracts from bovine tissues provide nutrients and support to the corresponding tissues in humans
- Vitamins, minerals, and nutrients from plants and animal tissues work synergistically for maximum effect†

#### Contains Protomorphogen™ extracts

- Standard Process' unique manufacturing method of deriving tissue cell determinants from animal glands and organs
- Help provide cellular support and rehabilitation in corresponding human tissues
- Important antigenic properties of nucleoprotein-mineral determinants, the foundation of the product†

### Unique Processing

#### Exclusive low-temperature, high-vacuum drying technique

- Preserves the enzymatic vitality and nutritional potential of ingredients

#### Degreed microbiologists and chemists in our on-site laboratories constantly conduct bacterial and analytical tests on raw materials, product batches, and finished products

- Ensures consistent quality and safety

#### Vitamin and mineral analyses validate product content and specifications

- Assures high-quality essential nutrients are delivered

### Whole Food Philosophy

Dr. Lee challenged common scientific beliefs by choosing a holistic approach of providing nutrients through whole foods. His goal was to provide nutrients as they are found in nature—in a whole food state where he believed their natural potency and efficacy would be realized. Dr. Lee believed that when nutrients remain intact and are not split from their natural associated synergists—known and unknown—bioactivity is markedly enhanced over synthetic nutrients. Following this philosophy, even a small amount of a whole food concentrate will offer enhanced nutritional support, compared to a synthetic or fractionated vitamin. Therefore, one should examine the source of nutrients rather than looking at the quantities of individual nutrients on product labels.

**Proprietary Blend:** *Tillandsia usneoides*, soy (bean), bovine liver PMG™ extract, betaine hydrochloride, calcium lactate, defatted wheat (germ), potassium bicarbonate, choline bitartrate, inositol, disodium phosphate, bovine adrenal Cytosol™ extract, oat flour, and ascorbic acid.

**Other Ingredients:** Gelatin, water, niacinamide, pyridoxal 5'-phosphate, colors, and calcium stearate.

**Suggested Use:** One capsule per meal, or as directed.

**Sold to health care professionals.**

Studies on nutrients generally use large doses and these studies, some of which are cited below, are the basis for much of the information we provide you in this publication about whole food ingredients. See the supplement facts for Betacol®.

Anderson L.E. 1998. *Mosby's Medical, Nursing, & Allied Health Dictionary*. 5th ed. St. Louis, MO: Mosby: 1366.  
Army N.P. Spanish Moss and Ball Moss. University of Florida: Cooperative Extension Service Institute of Food and Agricultural Sciences. Online. 21 May 2000.  
Balch J.F., Balch Phyllis A. 1997. *Prescription for Nutritional Healing*. 2nd ed. Garden City Park, NY: Avery Publishing Group: 15-16.  
Berdanier C.D. 1995. *Advanced Nutrition Micronutrients*. Boca Raton, FL: CRC Press: 94-105.  
Coffee C.J. 1998. *Metabolis*. 1st ed. Madison, CT: Fence Creek Publishing: 69.  
Costa M., Di Stasi L.C., Kirizawa M., et al. 1989. Screening in mice of some medicinal plants used for analgesic purposes in the state of Sao Paulo, part II. *J Ethnopharmacol*. 27(1-2): 25-33.  
Di Padova C., Bosisio E., Cighetti G., et al. 1982. 3-Hydroxy-3-methylglutaric acid (HMG) reduces dietary cholesterol induction of saturated bile in hamster. *Life Sci*. 30(22): 1907-1914.  
Duke J. USDA - ARS - NGR. Phytochemical and Ethnobotanical Database. Beltsville, MD: Beltsville Agricultural Research Center. Online. 22 May 2000.  
Guyton A.C., Hall J.E. 1997. *Human Physiology and Mechanisms of Disease*. 6th ed. Philadelphia, PA: W.B. Saunders Company: 589.  
Kirschmann J.D. 1979. *Nutrition Almanac*. Revised ed. New York, NY: McGraw-Hill Book Company: 25-27.  
Labate M.E., Dam R. 1980. Effect of 3-hydroxy-3-methylglutaric acid on cholesterol metabolism in female Japanese quail. *Poult Sci* 59(2): 383-389.  
Lupien P.J., Moorjani S., Brun D., et al. 1979. Effects of 3-hydroxy-3-methylglutaric acid on plasma and low-density lipoprotein cholesterol levels in familial hypercholesterolemia. *J Clin Pharmacol*. 19(2-3): 120-126.

Moorjani S., Lupien P.J. 1977. Effect in vitro of 3-hydroxy-3-methylglutaric acid on the synthesis of mevalonate and its precursors. *Arch Int Physiol Biochim* 85(1): 1-10.  
Murakami T., Nagamura Y., Hirano K. 1998. The recovering effect of betaine on carbon tetrachloride-induced liver injury. *J Nutr Sci Vitaminol* 44(2): 249-255.  
Pitchford P. 1993. *Healing With Whole Foods*. Revised ed. Berkeley, CA: North Atlantic Books: 122, 402-403.  
Savoie L.L., Lupien P.J. 1975. Organ distribution of 3-hydroxy-3-methylglutaric acid, a potential anticholesterolemic agent. *Can J Physiol Pharmacol* 53(4): 638-643.  
Savoie L.L., Lupien P.J. 1975. Preliminary toxicological investigations of 3-hydroxy-3-methylglutaric acid (HMG). I: Study of acute toxicity and of teratogenic activity in rats and mice. *Arzneimittelforschung* 25(8): 1284-1286.  
Shils M.E., Young V.R. 1988. *Modern Nutrition in Health and Disease*. 7th ed. Philadelphia, PA: Lea and Febiger: 370-381.  
*Tillandsia usneoides*. Ecology and Evolutionary Biology Conservatory. Online. 17 May 2000.  
Tver D.F., Russell P. 1989. *The Nutrition and Health Encyclopedia*. 2nd ed. New York, NY: Van Nostrand Reinhold: 366-368, 445-446.  
Van Niekerk J.L., Hendriks T., Gevers-Luven, J.A., et al. 1984. The lipid-lowering effects of 3-hydroxy-3-methylglutaric acid and bile acid drainage in WHHL rabbits. *Clin Sci* 67(4): 439-444.  
Wilson E.D., Fisher K.H., Fuqua M.E. 1965. *Principles of Nutrition*. 2nd ed. New York, NY: John Wiley & Sons, Inc: 290-294.  
Witherup K.M., McLaughlin J.L., Judd R.L., et al. 1995. Identification of 3-hydroxy-3-methylglutaric acid (HMG) as a hypoglycemic principle of Spanish moss (*Tillandsia usneoides*). *J Nat Prod* 58(8): 1285-1290.  
Yousufzai S.Y., Siddiqi M. 1976. 3-Hydroxy-3-methylglutaric acid and triton-induced hyperlipidemia in rats. *Experientia* 32(9): 1178-1179.  
Yousufzai S.Y., Siddiqi M. 1977. Serum and liver lipid responses to 3-hydroxy-3-methylglutaric acid in rats on different carbohydrate diets. *Lipids* 12(3): 262-266.